


IMAGES IN CARDIOLOGY

Singleton-Merten Syndrome and Impaired Cardiac Function

Israel Valverde, MD,*† Eric Rosenthal, MD,* Aphrodite Tzifa, MD,*† Pravin Desai, MD,‡ Aaron Bell, MD,* Kuberan Pushparajah, MD,* Shakeel Qureshi, MD,* Philipp Beerbaum, MD,*† John Simpson, MD*

London and Ipswich, United Kingdom



From the *Department of Pediatric Cardiology, Evelina Children's Hospital, Guy's and St. Thomas Foundation Trust, London, United Kingdom; †Division of Imaging Sciences, King's College London, London, United Kingdom; and the ‡Child Health Department, Ipswich Hospital National Health Service Trust, Ipswich, United Kingdom. Manuscript received February 19, 2010, accepted February 25, 2010.

A 13-year-old girl presented to our unit because of recurrent syncopal episodes. Echocardiography (A and B, [Online Video 1](#)) demonstrated left ventricular (LV) hypertrophy (*), poor aortic valve excursion, and intense echogenicity extending to the ascending aorta (arrows). Three-dimensional echocardiography showed dyskinesia of the LV with poor ejection fraction of 21%. The 32-slice computed tomography scan revealed the extension of the calcification up to the transverse arch (C to F, [Online Video 2](#)). The calcification was clearly delineated by fluoroscopy (arrow), with normal coronary angiogram (G to I).

The patient underwent aortic root replacement with pulmonary autograft. One week post-operatively, 3-dimensional echocardiography demonstrated dramatically improved LV ejection fraction (50%) with no evidence of dyskinesia ([Online Video 3](#)).

A diagnosis of Singleton-Merten syndrome, a rare genetic disorder characterized by calcium deposition, had been made recently. To our knowledge, cardiovascular imaging findings have not been reported in the literature.